

EARLY PSYCHOSIS DECLARATION

An International Consensus Statement¹ about Early Intervention and Recovery for Young People with Early Psychosis

Jointly issued by the
World Health Organization and International Early Psychosis Association

Introduction

According to the World Health Organization's World Health Report 2001, schizophrenia and other forms of psychoses which affect young people represent a major public health problem. Worldwide, they rank as the third most disabling condition, (following quadriplegia and dementia and higher than blindness and paraplegia) and pose an enormous burden, both in terms of economic cost and of human suffering.

Yet, in spite of the availability of interventions that can reduce relapses by more than 50%, not all affected individuals have access to them, and when they do, it is not always in a timely and sustained way. Among the goals of care to these people, the identification of the illness and its treatment, *as early as possible*, represents a high priority.

Comprehensive programmes for the detection and treatment of early psychosis and in supporting the needs of young people with early psychosis carry the important function of promoting recovery, independence, equity and self-sufficiency and of facilitating uptake of social, educational and employment opportunities for those young people.

These programmes can be provided by individuals and teams with specialised skills, with a full range of primary health care services for every young person with early psychosis.

Prompt and effective interventions for young people with early psychosis, for their families, close friends and other carers represent a major element of respect of individuals' rights to citizenship and social inclusion.

¹ **Acknowledgement:** The UK Newcastle Early Psychosis Declaration (2002) was an idea conceived by IRIS (Initiative to Reduce the Impact of Schizophrenia) and further developed with the support of Rethink and the National Institute of Mental Health for England. On behalf of these UK organisations Dr. David Shiers and Dr. Jo Smith have subsequently worked closely with Professor Pat McGorry (International Early Psychosis Association) and Dr. Jose Bertolote (World Health Organisation) to modify the declaration for an international audience in the international consensus statement.

Vision: Fundamental objectives derived from this statement are to:

- Challenge stigmatising and discriminatory attitudes so that young people are not disadvantaged by their experiences and are truly included in their local communities.
- Generate optimism and expectations of positive outcomes and recovery so that all young people with psychosis and their families achieve ordinary lives.
- Raise wider societal awareness about psychosis and the importance of early intervention.
- Attract and encourage practitioners from a wide range of health, social, non-governmental agencies (e.g. charitable, voluntary and youth), educational and employment services to reflect on how they can better contribute to supporting young people with psychosis, their families and their friends.

Values: Programmes for the detection and treatment of early psychosis should value:

- Respect of the right to recovery and social inclusion and support to the importance of personal, social, educational and employment outcomes.
- Respect of the strengths and qualities of young people with a psychosis, their families and communities, encouraging ordinary lives and expectations.
- Services that actively partner young people, their families and friends to place them at the centre of care and service delivery, at the same time sensitive to age, phase of illness, gender, sexuality and cultural background.
- Use of cost-effective interventions.
- Respect of the right for family and friends to participate and feel fully involved.

A five year programme of action

This consensus statement describes (table 1) a number of five year measurable outcomes that an individual and their family can expect from services who have successfully implemented a comprehensive and effective programme.

TABLE 1.
5 year outcomes which a comprehensive and effective programme would deliver to people with early psychosis and their families

Comprehensive programme	Measured Outcome
<p>Improving access, engagement and treatment:</p> <ul style="list-style-type: none"> ▪ Walk in responsive services usually provided in primary care settings should be equipped to deal effectively with early psychosis. ▪ Service interfaces are designed to support quicker and more effective engagements of young people 	<ul style="list-style-type: none"> ▪ The mean duration of untreated psychosis from the onset of psychosis (DuP) is less than three months ▪ The use of involuntary treatments in the first engagement is less than 25% ▪ Effective treatment will be provided after no more than 3 attempts to seek help
<p>Raising community awareness</p> <ul style="list-style-type: none"> ▪ about the importance and the opportunities for earlier detection and improved management of psychosis. 	<ul style="list-style-type: none"> ▪ All 15 year olds are equipped by mainstream education to understand and deal with psychosis ▪ Psychosis specific training is available to teachers and other relevant community agencies
<p>Promoting recovery</p> <ul style="list-style-type: none"> ▪ Services should enable rather than disable, ultimately aspiring to healing and recovery ▪ Means receiving encouragement and sufficient support to retain/regain full participation in society. 	<ul style="list-style-type: none"> ▪ Suicide rates within the first two years from diagnosis will be less than 1% ▪ Two years after diagnosis 90% of affected individuals have employment / education rates similar to their age / gender matched peers ▪ Two years after diagnosis 90% of affected individuals will report satisfaction with their employment, educational and social attainments ▪ Days out of role over the 2 years post diagnosis no more than 25%
<p>Family engagement and support</p> <ul style="list-style-type: none"> ▪ Families have better access to information and education, social, economic, practical and emotional support. 	<ul style="list-style-type: none"> ▪ Services will give a meaningful response to families or key supporters within one week ▪ 90% of families will feel respected and valued as partners in care
<p>Practitioner training:</p> <ul style="list-style-type: none"> ▪ All Primary care sites are equipped to deal effectively with early psychosis. ▪ Continued Professional Development is supported for all specialist staff working with young people with psychosis 	<ul style="list-style-type: none"> ▪ Recognition, care and treatment of young people with psychosis is a routine part of training curricula of all primary care and social care practitioners ▪ Specific Early Intervention training programmes are resourced and evaluated

Technical interventions: Attaining these 5 year outcomes will require underpinning by modern technical interventions and actions:

- A series of interventions has demonstrated efficacy in the management of early psychoses, among which there are pharmacological and psychosocial approaches, the latter addressed to both patients and family members. [refer International Early Psychosis Association consensus Clinical Practice Guidelines 2002]
- Some of these interventions are specified, by level of resources in a given area, in Table 2, within the framework provided (p.114 table 5.1) in the WHO's World Health Report (2001). The framework also suggests strategic actions necessary for the implementation of those interventions.

TABLE 2

Recommended minimum 5 year strategic actions required for the care of people with early psychosis – UK highlighted in Scenario C

10 overall strategic recommendations	Scenario A: Low level of resources	Scenario B: Medium level of resources	Scenario C: High level of resources as in UK
1. Provide treatment in primary care: <i>early detection and access to comprehensive community-based mental health services</i>	Primary care health personnel can recognise and initiate treatment for early psychosis	Pilot care pathway audits of those developing early psychosis to improve detection and treatment	<p>Develop strategies which reduce the time taken to access support from the onset of psychosis (Duration of Untreated Psychosis)</p> <p>Integrated Care Pathways are routinely evaluated based on agreed Referral Guidelines between primary care, relevant community agencies, and specialist services.</p> <p>Develop special access channels for these clients.</p>
2. Make psychotropic and psychosocial interventions available:	<p>Low dose typical antipsychotic and other relevant medications are available and offered routinely to people with early psychosis in all health care settings</p> <p>Practical psychosocial support available via family, friends and community organizers.</p>	<p>Initiate pilot mental health programmes to provide comprehensive care and treatment (<i>i.e psychotropic and psychosocial – the latter available to consumer and family</i>)</p> <p>Low dose atypical antipsychotic medication are available and offered routinely in all health care settings</p>	<p>Low dose atypical antipsychotic medicines are offered as drug treatment of first choice</p> <p>Cognitive behavioural therapy is routinely available</p> <p>Family interventions are routinely available</p> <p>The following treatment programmes are routinely provided: vocational recovery; continuing care; relapse prevention; substance use harm reduction.</p>
3. Give care in the community: <i>move towards least restrictive and least coercive service</i>	Move away from institutional and custodial models of care for those with early psychosis who have a persistent and disabling course and provide community care wherever possible for early	<p>Initiate schemes which divert people with early psychosis presenting to criminal justice services, more appropriately into mental health services</p> <p>Use of involuntary treatment in early</p>	<p>Develop strategies to minimise the use of involuntary treatments in early psychosis, e.g. home-based intervention, alternative residential settings.</p> <p>'Streamed' youth-focused in-patient services are routinely available</p> <p>Develop youth-friendly day programmes</p>

approaches	<p>psychosis</p> <p>Preferred in-patient facility of a general hospital psychiatric unit rather than traditional mental asylum</p>	<p>psychosis is measured routinely.</p>	<p>Develop youth oriented residential facilities to support independent living for young people with psychoses</p> <p>Strategies for reducing risk of suicide for people with early psychosis are developed</p> <p>Individuals with early psychosis are offered treatment and care in community settings wherever possible</p>
<p>4. Educate the public: raise community awareness about the importance and the opportunities for earlier detection and improved management of psychosis</p>	<p>Support public campaigns to raise awareness about mental illness in young people</p> <p>Support nongovernmental organizations in public education</p>	<p>Support campaigns directed at young people which reduce stigma and discrimination against mental illness</p> <p>Initiate evidence-based mental health promotion and early intervention programmes in collaboration with other sectors</p>	<p>Launch public campaigns to raise awareness about the recognition and treatment of early psychosis.</p> <p>As part of mainstream education all young people are taught about psychosis</p> <p>Psychosis awareness training is provided to teachers, youth and welfare workers, police and criminal justice services and other relevant community agencies</p>
<p>5. Involve communities, families and consumers:</p>	<p>Support the formation of self-help groups for young people with early psychosis, family members and key supporters</p> <p>Fund schemes for nongovernmental organizations and mental health initiatives</p>	<p>Pilot initiatives which help families and key supporters access help and information about early psychosis</p> <p>Representation of communities, consumers and families in services and policy making</p>	<p>Families and key supporters will receive effective services appropriate to their needs, and access to advocacy support as appropriate. Services should systematically demonstrate effectiveness in terms of</p> <ul style="list-style-type: none"> ▪ Early engagement of families and close friends ▪ whether families feel valued as partners in care
<p>6. Establish national policies, programmes and legislation</p>	<p>Formulate mental health programmes and policy</p> <p>Increase the budget for mental health care to at least 5% of GDP</p>	<p>Emerging policy and a financial commitment to improve services for people with early psychosis and their families</p> <p>Growth in mental health expenditure focused on young people with early psychosis.</p>	<p>Consumers and families are represented in service development, training and policy-making</p> <p>Early intervention in psychosis is embraced as a key national priority demonstrated by investment of resources</p> <p>Specific commitment in terms of policy and health spending which underpin services for young people with psychosis and their families</p>
<p>7. Develop human resources</p>	<p>Primary care health practitioners recognise, through awareness training, that young people with psychosis have their own legitimate mental health concerns and needs</p> <p>Awareness training of psychiatrists and</p>	<p>Specialist mental health practitioners receive specific training about the recognition, care and treatment of young people with early psychosis</p>	<p>Specialist mental health practitioners develop advanced skills and knowledge about early detection, care and treatment of psychotic disorders in young people.</p> <p>Recognition, care and treatment of young people with psychosis and their families should form a routine part of the training curricula of all primary (generalist) health and social care practitioners</p>

	nurses about early psychosis care and treatment		Psychosis awareness training is provided to teachers, youth and welfare workers, police and criminal justice services and other relevant community agencies
8. Link with other sectors: to facilitate the increasingly regarded concept of recovery	<p>Initiate school and workplace mental health programmes</p> <p>Encourage the activities of non-governmental organizations</p>	<p>Strengthen school and workplace mental health programmes</p> <p>Links are developed between substance misuse and psychosis care and treatment programmes</p>	<p>Develop effective youth-friendly integrated programmes for supporting access to work, education and recreation. Effectiveness should be demonstrated e.g.</p> <ul style="list-style-type: none"> ▪ client satisfaction ▪ employment rates ▪ educational achievement ▪ access to accommodation <p>Develop integrated programmes for care and treatment of those with a combination of early psychosis and substance misuse problems with a single unified service system.</p>
9. Monitor community mental health: relevant indicators of early psychosis	<p>Include mental disorders in basic health information systems</p> <p>Survey high-risk population groups</p>	Institute surveillance for early psychosis in the community	<p>Develop advanced mental health monitoring systems for early psychosis which can measure Duration of Untreated Psychosis; positive recovery outcomes; morbidity in health and social terms; mortality (including suicide); cost and burden</p> <p>Monitor effectiveness and cost-effectiveness of prevention and treatment programmes in early psychosis.</p>
10. Support more research	Conduct studies on the prevalence, course, outcome and impact of early psychosis in the community	Institute effectiveness and cost-effectiveness studies for management of early psychosis	<ul style="list-style-type: none"> ▪ on causes of psychosis ▪ on prevention of psychosis ▪ on novel psychosocial and psychopharmacological interventions ▪ on service delivery models <i>e.g. urban / rural / dispersed rural</i> <p>about prevalence and needs of children and adolescents with early psychosis (aged under 16) both in terms of early detection and intervention implications</p>

References:

International Early Psychosis Association **Draft Clinical Practice Guidelines on the Management of Early Psychosis, Ch 9** in: Edwards J. and McGorry P. Implementing Early Psychosis Services. *Dunitz: London, 2002.*

The World Health Report: 2001: **Mental health: new understanding, new hope.**
World Health Organisation: Geneva, 2001.